

# Prediabetes

## What is prediabetes?

Prediabetes is when your blood sugar (or glucose) levels are higher than normal but not high enough to be diagnosed as type 2 diabetes. Did you know that an estimated 57 million American adults—or 1 in every 4 adults—have prediabetes?

## What's going on in your body?

The beta cells of the body play an important role. The beta cells are in the pancreas, a large gland behind the stomach. These cells make and release insulin. Insulin is a hormone that is naturally made by the body. Insulin helps sugar move from your blood into your body's cells. Your cells need sugar for energy. In people with prediabetes:

- The beta cells may not be working as well as they should, and some may start to die off
- As the number of beta cells goes down, the pancreas may begin to make less insulin

This increases the risk of developing type 2 diabetes. But it *can* be prevented. See the box on this page to find out how.

## Can I reverse prediabetes?

**Yes! It may be possible to stop prediabetes and keep it from going on to type 2 diabetes. Eating a healthy diet and losing weight can help.** Losing just 5% to 10% of your body weight can make a difference. The best way to lose weight is with:

- A healthy eating plan
- Regular physical activity, such as walking, for 30 minutes a day, 5 days a week

## Who is at risk?

You are more likely to get prediabetes if you:

- Are overweight
- Are age 45 years or older
- Have a parent or sibling with diabetes
- Have high blood pressure
- Do not get regular physical activity

## How is it diagnosed?

You may have no symptoms. Instead, it is found with one of the following tests:

- Fasting glucose test (FGT)—Measures blood sugar when you haven't eaten anything for at least 8 hours
- Glucose tolerance test (GTT)—Measures blood sugar after you haven't eaten anything for at least 8 hours and 2 hours after you drink a sugary drink provided by a doctor or laboratory
- A1C—Measures your average estimated blood sugar over the past 2 to 3 months

Here are the results that the tests may show:

	Normal Blood Sugar	Prediabetes	Diabetes
FGT	Less than 100 mg/dL	100-125 mg/dL	126 mg/dL or higher
GTT	Less than 140 mg/dL	140-199 mg/dL	200 mg/dL or higher
A1C	Less than 5.7%	5.7%-6.4%	6.5% or higher

Adapted from the American Diabetes Association. Standards of medical care in diabetes—2011. *Diabetes Care*. 2011;34(suppl 1):S11-S61.

If results are normal, the tests should be repeated at least every 3 years.

For more information, visit [Cornerstones4Care.com](http://Cornerstones4Care.com)

Novo Nordisk Inc. grants permission to reproduce this piece for nonprofit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

Cornerstones4Care™ is a trademark of Novo Nordisk A/S.